

University of Groningen

Copper-catalyzed regio- and enantioselective synthesis of chiral enol acetates and beta-substituted aldehydes

Fananas-Mastral, M.; Feringa, B.L.

Published in:
Journal of the American Chemical Society

DOI:
[10.1021/ja105585y](https://doi.org/10.1021/ja105585y)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Fananas-Mastral, M., & Feringa, B. L. (2010). Copper-catalyzed regio- and enantioselective synthesis of chiral enol acetates and beta-substituted aldehydes. *Journal of the American Chemical Society*, 132(38), 13152 - 13153. <https://doi.org/10.1021/ja105585y>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SUPPORTING INFORMATION

Copper-Catalyzed Regio- and Enantioselective Synthesis of Chiral Enol Acetates and β -Substituted Aldehydes.

Martín Fañanás-Mastral and Ben L. Feringa*

*Department of Organic Chemistry, Stratingh Institute, University of Groningen, Nijenborgh 4,
9747 AG, Groningen, The Netherlands.*

General Procedures:

Chromatography: Merck silica gel type 9385 230-400 mesh, TLC: Merck silica gel 60, 0.25 mm. Components were visualized by UV and cerium/molibdenum staining. Progress and conversion of the reaction were determined by GC-MS (GC, HP6890: MS HP5973) with an HP1 or HP5 column (Agilent Technologies, Palo Alto, CA). Mass spectra were recorded on a AEI-MS-902 mass spectrometer (EI+) or a LTQ Orbitrap XL (ESI+). ^1H - and ^{13}C -NMR were recorded on a Varian AMX400 (400 and 100.59 MHz, respectively) or a Varian VXR300 (300 and 75 MHz, respectively) using CDCl_3 as solvent. Chemical shift values are reported in ppm with the solvent resonance as the internal standard (CHCl_3 : δ 7.26 for ^1H , δ 77.0 for ^{13}C). Data are reported as follows: chemical shifts, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants (Hz), and integration. Carbon assignments are based on APT ^{13}C -NMR experiments. Optical rotations were measured on a *Schmidt + Haensch* polarimeter (Polartronic MH8) with a 10 cm cell (*c* given in g/100 mL). Enantioselectivities were determined by HPLC analysis using a Shimadzu LC-10ADVP HPLC equipped with a Shimadzu SPD-M10AVP diode array detector or by capillary GC analysis (HP 6890, Chiraldex G-TA column (30 m x 0.25 mm)) using flame ionization detector.

All reactions were carried out under a nitrogen atmosphere using oven dried glassware and using standard Schlenk techniques. CH_2Cl_2 was dried and distilled over calcium hydride. Acetyl chloride, α,β -unsaturated aldehydes (**1**) and all copper-salts (CuI , CuTC , $\text{CuBr}\cdot\text{SMe}_2$) were purchased from Aldrich, and used without further purification. Grignard reagents were purchased from Aldrich (MeMgBr , EtMgBr , $n\text{-HexMgBr}$, $i\text{-BuMgBr}$). Grignard reagents were titrated using *s*-BuOH and catalytic amounts of 1,10-phenanthroline. Phosphoramidite ligands **L1-L3**,¹ **L4-L5**,² **L6**,³ **L7**,⁴ **L8**,⁵ **L9**,⁶ **L10**⁷ were prepared as reported in the literature.

Racemic products were synthesized by reaction of the α,β -unsaturated aldehydes (**1**) with acetyl chloride and ZnCl_2 and the corresponding Grignard reagent at -78°C in CH_2Cl_2 in the presence of CuI (10 mol%) and PPh_3 (12 mol%).

¹ Feringa, B. L.; Pineschi, M.; Arnold, L. A.; Imbos, R.; De Vries, A. H. M. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2620–2623.

² Tissot-Croset, K.; Polet, D.; Gille, S.; Hawner, C.; Alexakis, A. *Synthesis* **2004**, 2586-2590.

³ Teichert, J. F.; Feringa, B. L. *Synthesis*, **2010**, 1200-1204.


⁴ Arnold, L. A.; Imbos, R.; Mandoli, A.; De Vries, A. H. M.; Naasz, R.; Feringa, B. L. *Tetrahedron* **2000**, *56*, 2865-2878.

⁵ Fernández-Ibáñez, M. A.; Maciá, B.; Pizzuti, M. G.; Minnaard, A. J.; Feringa, B. L. *Angew. Chem. Int. Ed.* **2009**, *48*, 9339-9341.

⁶ Alexakis, A.; Polet, D.; Rosset, S.; March, S. *J. Org. Chem.* **2004**, *69*, 5660-5667.

⁷ Alexakis, A.; Burton, J.; Vastra, J.; Benhaim, C.; Fournioux, X.; van den Heuvel, A.; Levêque, J.-M.; Mazé, F.; Rosset, S. *Eur. J. Org. Chem.* **2000**, 4011 – 4028.

Table S1. Some copper sources, ligands and conditions tried in the optimization process



entry ^a	L	2a (%) ^b	3 (%) ^b	4 (%) ^b	Z/E	2a ee (%) ^f
1	L'1	12	35	10	1:2	Z: rac. E: rac.
2	L'2	24	32	14	1:2	Z: 42 E: 25
3	L'3	24	26	16	1:3	Z: rac. E: 4
4	L1 ^c	57	19	12	4:1	Z: 56 E: 78
5	L1 ^d	59	17	13	4:1	Z: 53 E: 51
6	L1 ^e	64	14	6	3:1	Z: 57 E: 56

^aEtMgBr added dropwise during 5 minutes. ^bIsolated yield. ^cCuTC:L4 5:5.5. ^dCuTC:L4 5:7.5. ^eCuTC:L4 5:10. ^fDetermined by HPLC analysis.






Table S2. Screening of different phosphoramidites.^a



entry	L	2a (%) ^b	3 (%) ^b	4 (%) ^b	Z/E ^c	2a ee (%) ^d	5a ee (%) ^e
1	L7	73	4	3	1.5:1	Z: 54 E: 55	55
2	L8	78	9	2	3:1	Z: -59 E: -75	-64 ^f
3	L9	52	10	7	4.5:1	Z: 38 E: 77	45
4	L10	53	12	-	3:1	Z: -6 E: -2	-5 ^f

^aReactions run on a 0.5 mmol scale adding 1.2 eq. of EtMgBr diluted with CH₂Cl₂ (0.7 mL) over 1h. ^bIsolated yield. ^cDetermined from the ¹H NMR spectrum. ^dDetermined by chiral HPLC. ^eDetermined by chiral GC. ^fNegative ee value indicates that the opposite enantiomer was formed.




General procedure for the synthesis of enol acetates **2 and transformation to the β -substituted aldehydes **5**.**

Copper thiophene carboxylate (CuTC) (0.025 mmol, 4.75 mg) and ligand **L1** (0.0275 mmol, 14.84 mg) were charged in a Schlenk tube equipped with septum and stirring bar. CH₂Cl₂ (2 mL) was added and the solution was stirred under nitrogen at room temperature for 30 min. In the meantime, the corresponding α,β -unsaturated aldehyde (0.5 mmol) was added dropwise to a solution of acetyl chloride (0.5 mmol, 38 μ L) and freshly fused ZnCl₂ (0.0075 mmol, 1 mg) in CH₂Cl₂ (1 mL) at -10°C. Then, the resulting solution was added to the catalyst solution which was previously cooled to -78°C. After 5 min. at this temperature the corresponding Grignard reagent (solution in Et₂O, 0.6 mmol) was diluted with CH₂Cl₂ (0.7 mL) and was added dropwise over 6 hours using a syringe pump. Once the addition was complete, the mixture was stirred other four hours at -78°C. The reaction was quenched with aqueous saturated NH₄Cl (2 mL) and the mixture was warmed up to room temperature. The mixture was diluted with Et₂O and the layers were separated. The aqueous layer was extracted with Et₂O (3 x 5 mL) and the combined organic layers were dried with anhydrous Na₂SO₄, filtered and the solvent was evaporated *in vacuo*. At this point GC analysis was carried out to determine the b:l ratio. The crude product was purified by flash chromatography on silica gel using different mixtures of Pentane:Et₂O as eluents.


Hydrolysis of the enol acetate was then performed using the following procedure: enol acetate **2** was dissolved in MeOH (10 mL) and K₂CO₃ (5 eq.) was added. The mixture was stirred at room temperature during one hour. Then the solvent was removed *in vacuo*, water (5 mL) was added and the mixture was extracted with Et₂O (3 x 5 mL). The combined organic layers were dried with anhydrous Na₂SO₄, filtered and the solvent was evaporated *in vacuo* to afford the corresponding aldehyde **5**.

*One-pot process for the synthesis of β -substituted aldehydes **5**.*

After the reaction was complete, it was quenched with MeOH (5 mL) and was warmed up to room temperature. Then K₂CO₃ (5 eq.) was added and the mixture was stirred for one hour. Same work-up gave rise to the aldehyde **5**.




(S)-3-phenylpent-1-enyl acetate (2a): Colorless oil obtained as a 13:1 mixture of *Z* (major) and *E* (minor) isomers in 89% yield after column chromatography using a mixture of Pentane:Et₂O 40:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.33-7.29 (m, 2H), 7.26-7.19 (m, 3H), 7.11 (d, *J* = 6.4, 1H), 5.04 (dd, *J* = 9.8, 6.4, 1H), 3.73 (q, *J* = 8.3, 1H), 2.16 (s, 3H), 1.82-1.65 (m, 2H), 0.90 (t, *J* = 7.3, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.2, 144.8, 133.9, 128.7, 127.5, 126.4, 117.7, 43.0, 29.6, 21.0, 12.3. HRMS (ESI+, *m/z*): calcd for C₁₃H₁₆O₂Na [M+Na]⁺: 227.10425; found: 227.10400. *Ee* was determined by chiral HPLC analysis, Chiralpak OJ-H column, Heptane/*i*-PrOH 95:5, retention times: 12.6 (*Z*, minor enantiomer), 14.0 (*E*, major enantiomer), 16.4 (*E*, minor enantiomer), and 20.4 (*Z*, major enantiomer) min. [*Z* isomer: 91% ee; *E* isomer: 92% ee].




(S)-3-phenylnon-1-enyl acetate (2b): Colorless oil obtained as a 10:1 mixture of *Z* (major) and *E* (minor) isomers in 88% yield after column chromatography using a mixture of Pentane:Et₂O 50:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.33-7.29 (m, 2H), 7.25-7.19 (m, 3H), 7.10 (d, *J* = 6.4, 1H), 5.04 (dd, *J* = 9.8, 6.4, 1H), 3.73 (q, *J* = 8.2, 1H), 2.16 (s, 3H), 1.79-1.61 (m, 2H), 1.37-1.19 (m, 8H), 0.89 (t, *J* = 6.6, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.2, 145.0, 133.7, 128.7, 127.5, 126.4, 118.0, 41.2, 36.7, 32.0, 29.4, 27.6, 22.9, 21.0, 14.3. HRMS (ESI+, *m/z*): calcd for C₁₇H₂₄O₂Na [M+Na]⁺: 283.16685; found: 283.16528.

Ee was determined by transformation into the aldehyde **5b** following the general procedure (94% ee, *vide infra*).




(S)-3-phenylbut-1-enyl acetate (2c): Colorless oil obtained as a 7:1 mixture of *Z* (major) and *E* (minor) isomers in 66% yield after column chromatography using a mixture of Pentane:Et₂O 40:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.34-7.22 (m, 5H), 7.07 (d, *J* = 6.3, 1H), 5.06 (dd, *J* = 9.5, 6.3, 1H), 4.03 (quint, *J* = 7.6, 1H), 2.17 (s, 3H), 1.39 (d, *J* = 7.2, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.2, 145.8, 133.1, 128.7, 127.0, 126.4, 119.1, 35.2, 22.0, 21.0. HRMS (ESI+, *m/z*): calcd for C₁₂H₁₄O₂Na [M+Na]⁺: 213.08860; found: 213.08682.

Ee was determined by chiral HPLC analysis, Chiralpak OJ-H column, Heptane/*i*-PrOH 95:5, retention times: 13.5 (*Z*, minor enantiomer), 15.2 (*E*, minor enantiomer), 17.4 (*Z*, major enantiomer), and 19.8 (*E*, major enantiomer) min. [*Z* isomer: 83% ee; *E* isomer: 4% ee].




(S)-5-methyl-3-phenylhex-1-enyl acetate (2d): Colorless oil obtained as a 12:1 mixture of *Z* (major) and *E* (minor) isomers in 85% yield after column chromatography using a mixture of Pentane:Et₂O 50:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.32-7.30 (m, 2H), 7.26-7.19 (m, 3H), 7.09 (d, *J* = 6.4, 1H), 5.02 (dd, *J* = 9.8, 6.4, 1H), 3.94 (q, *J* = 7.9, 1H), 2.17 (s, 3H), 1.63-1.49 (m, 3H), 0.95 (d, *J* = 6.5, 3H), 0.93 (d, *J* = 6.6, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.1, 145.1, 133.7, 128.7, 127.5, 126.4, 118.1, 46.1, 39.1, 25.9, 23.2, 22.4, 21.0. HRMS (ESI+, *m/z*): calcd for C₁₅H₂₀O₂Na [M+Na]⁺: 255.1356; found: 255.1353.

Ee was determined by chiral HPLC analysis, Chiralpak OJ-H column, Heptane/*i*-PrOH 99:1, retention times: 12.5 (*Z*, minor enantiomer), 14.5 (*E*, minor enantiomer), 16.2 (*Z*, major enantiomer), and 18.3 (*E*, major enantiomer) min. [*Z* isomer: 49% ee; *E* isomer: 45% ee].




(S)-3-(furan-2-yl)pent-1-enyl acetate (2e): In this case the corresponding chloroacetate was formed at -78°C during 5 minutes. Colorless oil obtained as a 18:1 mixture of *Z* (major) and *E* (minor) isomers in 75% yield after column chromatography using a mixture of Pentane:Et₂O 40:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.32 (br s, 1H), 7.16 (d, $J = 6.4$, 1H), 6.29 (br s, 1H), 6.02 (br s, 1H), 4.96 (dd, $J = 9.7$, 6.4, 1H), 3.85 (q, $J = 8.1$, 1H), 2.16 (s, 3H), 1.91-1.81 (m, 1H), 1.66- 1.55 (m, 1H), 0.90 (t, $J = 7.5$, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.0, 157.3, 146.9, 134.8, 114.4, 110.2, 104.7, 36.6, 27.2, 21.0, 11.7. HRMS (ESI+, m/z): calcd for C₁₁H₁₄O₃Na [M+Na]⁺: 217.0835; found: 217.0828.

Ee was determined by transformation into the aldehyde **5e** following the general procedure (90% *ee*, *vide infra*).




(S)-3-(furan-2-yl)non-1-enyl acetate (2f): In this case the corresponding chloroacetate was formed at -78°C during 5 minutes. Colorless oil obtained as a 17:1 mixture of *Z* (major) and *E* (minor) isomers in 69% yield after column chromatography using a mixture of Pentane:Et₂O 40:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.31 (br s, 1H), 7.14 (d, $J = 6.4$, 1H), 6.28 (br s, 1H), 6.01 (br s, 1H), 4.96 (dd, $J = 9.9$, 6.4, 1H), 3.85 (q, $J = 8.1$, 1H), 2.15 (s, 3H), 1.85-1.75 (m, 1H), 1.60- 1.52 (m, 1H), 1.33-1.21 (m, 8H), 0.88 (t, $J = 6.9$, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.1, 157.5, 141.2, 134.6, 114.7, 110.2, 104.5, 34.9, 34.1, 31.9, 29.3, 27.1, 22.9, 21.0, 14.3. HRMS (ESI+, m/z): calcd for C₁₅H₂₂O₃Na [M+Na]⁺: 273.1461; found: 273.1454.

Ee was determined by transformation into the aldehyde **5f** following the general procedure (90% *ee*, *vide infra*).




(S)-3-(2-methoxyphenyl)pent-1-enyl acetate (2g): In this case the corresponding chloroacetate was formed at -78°C during 5 minutes. Colorless oil obtained as a 16:1 mixture of *Z* (major) and *E* (minor) isomers in 77% yield after column chromatography using a mixture of Pentane:Et₂O 30:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.20-7.17 (m, 2H), 7.06 (d, *J* = 6.4, 1H), 6.92 (t, *J* = 7.4, 1H), 6.87 (d, *J* = 8.3, 1H), 5.15 (dd, *J* = 9.8, 6.4, 1H), 4.09 (q, *J* = 8.2, 1H), 3.83 (s, 3H), 2.14 (s, 3H), 1.80-1.62 (m, 2H), 0.88 (t, *J* = 7.4, 4H). ¹³C NMR (100 MHz, CDCl₃) δ 168.4, 157.3, 133.8, 133.0, 128.2, 127.3, 120.8, 117.4, 111.0, 55.6, 37.1, 28.5, 21.0, 12.4. HRMS (ESI+, *m/z*): calcd for C₁₄H₁₈O₃Na [M+Na]⁺: 257.11482; found: 257.11298.

Ee was determined by transformation into the aldehyde **5g** following the general procedure (91% *ee*, *vide infra*).




(S)-3-(2-methoxyphenyl)non-1-enyl acetate (2h): In this case the corresponding chloroacetate was formed at -78°C during 5 minutes. Colorless oil obtained as a 20:1 mixture of *Z* (major) and *E* (minor) isomers in 81% yield after column chromatography using a mixture of Pentane:Et₂O 50:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.19-7.16 (m, 2H), 7.03 (d, *J* = 6.4, 1H), 6.91 (t, *J* = 7.4, 1H), 6.86 (d, *J* = 8.5, 1H), 5.14 (dd, *J* = 9.7, 6.4, 1H), 4.17 (q, *J* = 8.2, 1H), 3.83 (s, 3H), 2.14 (s, 3H), 1.76-1.54 (m, 2H), 1.34-1.21 (m, 8H), 0.88 (t, *J* = 6.6, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.3, 157.2, 133.6, 133.3, 128.1, 127.2, 120.8, 117.7, 111.1, 55.7, 35.6, 35.3, 32.0, 29.4, 27.7, 22.9, 21.0, 14.3. HRMS (ESI+, *m/z*): calcd for C₁₈H₂₆O₃Na [M+Na]⁺: 313.17742; found: 313.17734.

Ee was determined by transformation into the aldehyde **5h** following the general procedure (93% *ee*, *vide infra*).




(S)-3-methylnon-1-enyl acetate (2i): Colorless oil obtained as a 2:1 mixture of *Z* (major) and *E* (minor) isomers in 86% yield after column chromatography using a mixture of Pentane:Et₂O 80:1 as eluent. For the *Z* isomer: ¹H NMR (400 MHz, CDCl₃) δ 6.96 (d, *J* = 6.4, 1H), 4.65 (dd, *J* = 9.6, 6.4, 1H), 2.68-2.60 (m, 1H), 2.12 (s, 3H), 1.30-1.20 (m, 10H), 0.96 (d, *J* = 6.6, 3H), 0.87 (t, *J* = 6.7, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.4, 133.1, 120.6, 37.4, 32.7, 32.1, 29.6, 27.5, 22.9, 21.2, 21.0, 14.3. For the *E* isomer: ¹H NMR (400 MHz, CDCl₃) δ 7.05 (d, *J* = 12.5, 1H), 5.28 (dd, *J* = 12.5, 8.7, 1H), 2.15-2.07 (m, 1H), 2.09 (s, 3H), 1.30-1.20 (m, 10H), 0.99 (d, *J* = 6.7, 3H), 0.87 (t, *J* = 6.7, 3H). ¹³C NMR (100 MHz, CDCl₃) δ 168.5, 134.6, 121.0, 37.4, 32.0, 29.7, 29.5, 27.4, 22.9, 21.2, 21.0, 14.3. HRMS (ESI+, *m/z*): calcd for C₁₂H₂₂O₂Na [M+Na]⁺: 221.15120; found: 221.15100. *Ee* was determined by transformation into the aldehyde **5i** following the general procedure (92% *ee*, *vide infra*).




(S)-3-phenylpentanal (5a): Colorless oil obtained in 89% yield; 92% *ee*. The physical data were identical in all respects to those previously reported.⁸ *Ee* was determined by chiral GC analysis, Chiraldex G-TA column (30 m x 0.25 mm), (90°C isotherm), retention times: 40.4 (major enantiomer) and 41.9 min. [α]_D²⁰ = +11.5 (*c* = 1.0, CHCl₃).


⁸ Bräse, S.; Höfener, S. *Angew. Chem., Int. Ed.* **2005**, *44*, 7879-7881.



(S)-3-phenylnonananal (5b): Colorless oil obtained in 88% yield; 94% *ee*. The physical data were identical in all respects to those previously reported for the racemic compound.⁹ *Ee* was determined by chiral GC analysis, Chiraldex G-TA column (30 m x 0.25 mm), (110°C isotherm), retention times: 46.9 (major enantiomer) and 47.2 min. $[\alpha]_D^{20} = +10.8$ ($c = 1.0$, CHCl_3).



(S)-3-phenylbutanal (5c): Colorless oil obtained in 66% yield; 72% *ee*. The physical data were identical in all respects to those previously reported.¹⁰ *Ee* was determined by chiral GC analysis, Chiraldex G-TA column (30 m x 0.25 mm), (110°C isotherm), retention times: 46.9 (major enantiomer) and 47.2 min. $[\alpha]_D^{20} = +18.4$ ($c = 0.25$, CHCl_3).




(S)-5-methyl-3-phenylhexanal (5d): Colorless oil obtained in 80% yield; 48% *ee*. ¹H NMR (400 MHz, CDCl_3) δ 9.65 (t, $J = 2.1$, 1H), 7.33-7.29 (m, 2H), 7.23-7.18 (m, 3H), 3.33-3.25 (m, 1H), 2.71 (ddd, $J = 16.5$, 7.7, 2.1, 1H), 2.65 (ddd, $J = 16.5$, 6.8, 2.2, 1H), 1.63 (ddd, $J = 13.1$, 10.1, 4.6, 1H), 1.47-1.32 (m, 2H), 0.91 (d, $J = 6.3$, 3H), 0.85 (d, $J = 6.5$, 3H). ¹³C NMR (100 MHz, CDCl_3) δ 202.0, 143.9, 128.6, 127.5, 126.5, 51.1, 45.8, 37.9, 25.2, 23.3, 21.7. HRMS (ESI+, m/z): calcd for $\text{C}_{13}\text{H}_{18}\text{ONa}$ $[\text{M}+\text{Na}]^+$: 213.1250; found: 213.1243.


Ee was determined by chiral GC analysis, Chiraldex G-TA column (30 m x 0.25 mm), (90°C isotherm), retention times: 45.5 (major enantiomer) and 45.9 min. $[\alpha]_D^{20} = +5.4$ ($c = 1.0$, CHCl_3).

⁹ Marshall, J. A.; Herold, M.; Eidam, H. S.; Eidam, P. *Org. Lett.* **2006**, 8, 5505-5508.


¹⁰ Bull, S. D.; Davies, S. G.; Nicholson, R. L.; Sanganeer, H. J.; Smith, A. D. *Org. Biomol. Chem.* **2003**, 1, 2886 - 2899.




(S)-3-(furan-2-yl)pentanal (5e): Pale yellow oil obtained in 68% yield; 90% *ee*. ^1H NMR (400 MHz, CDCl_3) δ 9.71 (t, $J = 1.6$, 1H), 7.30 (br s, 1H), 6.26 (br s, 1H), 6.02 (d, $J = 1.9$, 1H), 3.22 (quint., $J = 7.1$, 1H), 2.74 (ddd, $J = 16.6$, 7.7, 1.8, 1H), 2.63 (ddd, $J = 16.6$, 6.4, 1.3, 1H), 1.73-1.62 (m, 2H), 0.86 (t, $J = 7.3$, 3H). HRMS (ESI+, m/z): calcd for $\text{C}_9\text{H}_{13}\text{O}_2$ $[\text{M}+\text{H}]^+$: 153.09101; found: 153.08955. *Ee* was determined by chiral GC analysis, Chiraldex G-TA column (30 m x 0.25 mm), (90°C isotherm), retention times: 13.8 (major enantiomer) and 14.9 min. $[\alpha]_{\text{D}}^{20} = +30.4$ ($c = 0.5$, CHCl_3).



(S)-3-(furan-2-yl)nonanal (5f): Pale yellow oil obtained in 63% yield; 90% *ee*. ^1H NMR (300 MHz, CDCl_3) δ 9.71 (t, $J = 2.0$, 1H), 7.31 (d, $J = 1.0$, 1H), 6.27 (dd, $J = 3.1$, 1.8, 1H), 6.02 (d, $J = 3.1$, 1H), 3.33- 3.24 (m, 1H), 2.74 (ddd, $J = 16.7$, 7.8, 2.1, 1H), 2.64 (ddd, $J = 16.7$, 6.3, 1.9, 1H), 1.73-1.54 (m, 2H), 1.35-1.15 (m, 8H), 0.86 (t, $J = 6.8$, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 201.2, 154.9, 141.4, 109.8, 105.3, 47.7, 39.1, 33.5, 31.7, 29.1, 27.2, 22.6, 14.0. HRMS (ESI+, m/z): calcd for $\text{C}_{13}\text{H}_{21}\text{O}_2$ $[\text{M}+\text{H}]^+$: 209.15361; found: 209.15322. *Ee* was determined by chiral GC analysis, Chiraldex G-TA column (30 m x 0.25 mm), (110°C isotherm), retention times: 32.7 (major enantiomer) and 34.4 min. $[\alpha]_{\text{D}}^{20} = +4.8$ ($c = 0.92$, CHCl_3).




(S)-3-(2-methoxyphenyl)pentanal (5g): Colorless oil obtained in 77% yield; 91% *ee*. The physical data were identical in all respects to those previously reported.⁸ *Ee* was determined by chiral GC analysis, Chiraldex G-TA column (30 m x 0.25 mm), (110°C isotherm), retention times: 33.4 (major enantiomer) and 34.7 min. $[\alpha]_D^{20} = -19.3$ ($c = 2.7$, CHCl_3).



(S)-3-(2-methoxyphenyl)nonanal (5h): Colorless oil obtained in 79% yield; 93% *ee*. ^1H NMR (400 MHz, CDCl_3) δ 9.63 (t, $J = 2.0$, 1H), 7.19 (t, $J = 7.7$, 1H), 7.14 (d, $J = 7.4$, 1H), 6.92 (t, $J = 7.4$, 1H), 6.86 (d, $J = 8.1$, 1H), 3.81 (s, 3H), 3.66-3.57 (m, 1H), 6.86 (dd, $J = 7.1$, 2.0, 2H), 1.76-1.52 (m, 2H), 1.35-1.10 (m, 8H), 0.85 (t, $J = 6.4$, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 203.3, 157.4, 132.0, 128.0, 127.6, 120.9, 110.9, 55.5, 49.8, 35.1, 33.5, 31.9, 29.5, 27.5, 22.8, 14.3. HRMS (ESI+, m/z): calcd for $\text{C}_{16}\text{H}_{25}\text{O}_2$ $[\text{M}+\text{H}]^+$: 249.18491; found: 249.18463.


Ee was determined by chiral HPLC analysis, Chiralpak AD-H column, Heptane/*i*-PrOH 99.5:0.5, retention times: 12.2 (major enantiomer) and 12.8 min. $[\alpha]_D^{20} = -2.2$ ($c = 0.93$, CHCl_3).



(S)-3-methylnonan-1-ol (5i): Colorless oil obtained in 81% yield; 92% *ee*. The physical data were identical in all respects to those previously reported for the racemic compound.⁹ *Ee* was determined by chiral GC analysis, Chiraldex G-TA column (30 m x 0.25 mm), (60°C isotherm), retention times: 39.6 (major enantiomer) and 39.9 min. $[\alpha]_D^{20} = -18.4$ ($c = 1.95$, CHCl_3).

General procedure for the synthesis of aryl ketone 6.¹¹

To a solution of 2-bromonaphthalene (0.3 mmol, 62 mg) in anhydrous dimethylsulfoxide (0.6 mL) was added the enol acetate **2a** (0.6 mmol, 122 mg), tributyltin methoxide (0.6 mmol, 0.17 mL) followed by $\text{PdCl}_2[(o\text{-Tol})_3\text{P}]_2$ (0.015 mmol, 12 mg). The mixture was heated under nitrogen at 100°C for 16 hours and then cooled to room temperature. The mixture was diluted with EtOAc (10 mL) and a solution of aqueous KF 4M (5 mL) was added. The mixture was vigorously stirred over 1 hour and the filtered through celite. The organic layer was washed with water (3 x 5 mL) and dried with anhydrous Na_2SO_4 . The solvent was evaporated *in vacuo* and the product was purified by column chromatography on silica gel (Pentane : Et_2O 50:1).



(S)-1-(naphthalen-2-yl)-3-phenylpentan-1-one (6): Colorless oil obtained in 71% yield; 91% *ee*. ^1H NMR (400 MHz, CDCl_3) δ 8.42 (s, 1H), 8.00 (d, $J = 8.6$, 1H), 7.95 (d, $J = 8.0$, 1H), 7.87 (d, $J = 8.6$, 2H), 7.60 (t, $J = 7.9$, 1H), 7.55 (t, $J = 7.4$, 1H), 7.34-7.28 (m, 5H), 3.42 (t, $J = 6.9$, 2H), 3.38-3.29 (m, 1H), 1.92-1.82 (m, 1H), 1.77-1.66 (m, 1H), 0.86 (t, $J = 7.4$, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 199.4, 145.0, 135.7, 134.8, 132.7, 129.9,

¹¹ Jean, M.; Renault, J.; Uriac, P.; Capet, M.; van de Weghe, P. *Org. Lett.* **2007**, 9, 3623-3625.

129.8, 128.7, 128.6, 128.0, 127.9, 127.0, 126.5, 124.2, 46.0, 43.4, 29.5, 12.4. HRMS (APCI+, m/z): calcd for $C_{21}H_{21}O$ $[M+H]^+$: 289.1587; found: 289.1568.

Ee was determined by chiral HPLC analysis, Chiralpak AD-H column, Heptane/*i*-PrOH 95:5, retention times: 14.4 (major enantiomer) and 18.2 min. $[\alpha]_D^{20} = -23.3$ ($c = 1.2$, $CHCl_3$).